

chain nodes :

13

ring nodes :

1 2 3 4 5 6 7 8 9 10 11 12 14 15 16 17

chain bonds :

10-13

ring bonds :

1-2 1-6 2-3 3-4 4-5 4-7 4-16 5-6 5-9 5-15 7-8 7-14 7-10 8-9 10-11
11-12 12-14 15-17 16-17

exact/norm bonds :

1-2 1-6 2-3 3-4 4-5 4-7 4-16 5-6 5-9 5-15 7-8 7-14 7-10 8-9 10-13
10-11 11-12 12-14 15-17 16-17

Match level :


1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom
11:Atom 12:Atom 13:CLASS 14:Atom 15:Atom 16:Atom 17:Atom

L7 STRUCTURE UPLOADED

=> d

L7 HAS NO ANSWERS

L7 STR



Structure attributes must be viewed using STN Express query preparation.

=> s 17 full

FULL SEARCH INITIATED 20:11:18 FILE 'REGISTRY'

FULL SCREEN SEARCH COMPLETED - 1251 TO ITERATE

100.0% PROCESSED 1251 ITERATIONS

SEARCH TIME: 00.00.01


34 ANSWERS

L8 34 SEA SSS FUL L7

=> d 18 1-15

L8 ANSWER 1 OF 34 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 873078-37-6 REGISTRY
 ED Entered STN: 31 Jan 2006
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione,
 2,3-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1S,3'aS,5R,7'aS)-
 (9CI) (CA INDEX NAME)
 OTHER NAMES:
 CN Dechlorodauricumine
 FS STEREOSEARCH
 MF C19 H25 N O6
 SR CA
 LC STN Files: CA, CAPLUS

Absolute stereochemistry. Rotation (+).




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

1 REFERENCES IN FILE CA (1907 TO DATE)
 1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L8 ANSWER 2 OF 34 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 637770-99-1 REGISTRY
 ED Entered STN: 15 Jan 2004
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione,
 9'-chloro-2',3'-dihydro-5-hydroxy-6',7'-dimethoxy-1'-methyl-,
 (1R,3'aS,5R,7'aS,9'S)- (9CI) (CA INDEX NAME)
 FS STEREOSEARCH
 MF C18 H22 Cl N O5
 SR CA
 LC STN Files: CA, CAPLUS, USPATFULL

Absolute stereochemistry.




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

1 REFERENCES IN FILE CA (1907 TO DATE)
 1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L8 ANSWER 3 OF 34 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 637770-98-0 REGISTRY
 ED Entered STN: 15 Jan 2004
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,4',5'-trione,
 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-,
 (1R,3'aS,5S,7'aS,9'S)- (9CI) (CA INDEX NAME)
 FS STEREOSEARCH
 MF C19 H22 Cl N O7
 SR CA
 LC STN Files: CA, CAPLUS, USPATFULL

Absolute stereochemistry.




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

1 REFERENCES IN FILE CA (1907 TO DATE)
 1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L8 ANSWER 4 OF 34 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 637770-97-9 REGISTRY
 ED Entered STN: 15 Jan 2004
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione,
 9'-chloro-2',3'-dihydro-4,5,6',7'-tetramethoxy-1'-methyl-,
 (1R,3'aS,5S,7'aS,9'S)- (9CI) (CA INDEX NAME)
 FS STEREOSEARCH
 MF C20 H26 Cl N O6
 SR CA
 LC STN Files: CA, CAPLUS, USPATFULL

Absolute stereochemistry.




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

1 REFERENCES IN FILE CA (1907 TO DATE)
 1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L8 ANSWER 5 OF 34 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 637770-96-8 REGISTRY
 ED Entered STN: 15 Jan 2004
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione,
 9'-chloro-1'-ethyl-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-, 5'-oxime,
 (1R,3'aS,5S,7'aS,9'S)-(9CI) (CA INDEX NAME)
 FS STEREOSEARCH
 MF C20 H27 Cl N2 O6
 SR CA
 LC STN Files: CA, CAPLUS, USPATFULL

Absolute stereochemistry.
 Double bond geometry unknown.




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

1 REFERENCES IN FILE CA (1907 TO DATE)
 1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L8 ANSWER 6 OF 34 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 637770-95-7 REGISTRY
 ED Entered STN: 15 Jan 2004
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione,
 9'-chloro-2',3'-dihydro-4,6',7'-trimethoxy-5-(1-oxopropoxy)-1'-(1-oxopropoxy)-, (1R,3'aS,5S,7'aS,9'S)-(9CI) (CA INDEX NAME)
 FS STEREOSEARCH
 MF C24 H30 Cl N O8
 SR CA
 LC STN Files: CA, CAPLUS, USPATFULL

Absolute stereochemistry.




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

1 REFERENCES IN FILE CA (1907 TO DATE)
 1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L8 ANSWER 7 OF 34 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 637770-94-6 REGISTRY
 ED Entered STN: 15 Jan 2004
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione,
 9'-chloro-1'-ethyl-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-,
 (1R,3'aS,5S,7'aS,9'S)-(9CI) (CA INDEX NAME)
 FS STEREOSEARCH
 MF C20 H26 Cl N O6
 SR CA
 LC STN Files: CA, CAPLUS, USPATFULL

Absolute stereochemistry.




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

1 REFERENCES IN FILE CA (1907 TO DATE)
 1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L8 ANSWER 8 OF 34 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 637770-93-5 REGISTRY
 ED Entered STN: 15 Jan 2004
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione,
 9'-chloro-2',3'-dihydro-4,6',7'-trimethoxy-1'-methyl-5-(1-oxopropoxy)-,
 (1R,3'aS,5S,7'aS,9'S)-(9CI) (CA INDEX NAME)
 FS STEREOSEARCH
 MF C22 H28 Cl N O7
 SR CA
 LC STN Files: CA, CAPLUS, USPATFULL

Absolute stereochemistry.




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

1 REFERENCES IN FILE CA (1907 TO DATE)
 1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L8 ANSWER 9 OF 34 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 488736-04-5 REGISTRY
 ED Entered STN: 11 Feb 2003
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione,
 2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1S,3'aS,5R,7'aS)-
 (9CI) (CA INDEX NAME)
 FS STEREOSEARCH
 MF C19 H25 N O6
 SR CA
 LC STN Files: CA, CAPLUS, TOXCENTER

Absolute stereochemistry. Rotation (-).




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

1 REFERENCES IN FILE CA (1907 TO DATE)
 1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L8 ANSWER 10 OF 34 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 488736-03-4 REGISTRY
 ED Entered STN: 11 Feb 2003
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione,
 2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-, (1S,3'aS,5S,7'aS)- (9CI) (CA
 INDEX NAME)
 FS STEREOSEARCH
 MF C18 H23 N O6
 SR CA
 LC STN Files: CA, CAPLUS, TOXCENTER

Absolute stereochemistry. Rotation (-).




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

1 REFERENCES IN FILE CA (1907 TO DATE)
 1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L8 ANSWER 11 OF 34 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 486429-90-7 REGISTRY
 ED Entered STN: 06 Feb 2003
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione,
 9'-chloro-2',3'-dihydro-1-[3-(4-hydroxyphenyl)-2-nitropropyl]-4,6',7'-
 trimethoxy-, (1S,3'aS,5R,7'aS,9'S)- (9CI) (CA INDEX NAME)
 OTHER NAMES:
 CN Nitrotyrasacutuminine
 FS STEREOSEARCH
 MF C27 H31 Cl N2 O8
 SR CA
 LC STN Files: CA, CAPLUS

Absolute stereochemistry. Rotation (-).
 Currently available stereo shown.




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

1 REFERENCES IN FILE CA (1907 TO DATE)
 1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L8 ANSWER 12 OF 34 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 345641-00-1 REGISTRY
 ED Entered STN: 12 Jul 2001
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione,
 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1-methyl-,
 (1R,3'aS,5R,7'aS,9'S)- (CA INDEX NAME)
 OTHER NAMES:
 CN Dauricumine
 FS STEREOSEARCH
 MF C19 H24 Cl N O6
 SR CA
 LC STN Files: CA, CAPLUS, TOXCENTER

Absolute stereochemistry. Rotation (-).




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

3 REFERENCES IN FILE CA (1907 TO DATE)
 3 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L8 ANSWER 13 OF 34 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 345640-99-5 REGISTRY
 ED Entered STN: 12 Jul 2001
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione,
 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-,
 (1R,3'aS,5R,7'aS,9'S)-(9CI) (CA INDEX NAME)
 OTHER NAMES:
 CN Dauricumidine
 FS STEREOSEARCH
 MF C18 H22 Cl N O6
 SR CA
 LC STN Files: CA, CAPLUS

Absolute stereochemistry. Rotation (+).




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

2 REFERENCES IN FILE CA (1907 TO DATE)
 2 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L8 ANSWER 14 OF 34 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 219794-33-9 REGISTRY
 ED Entered STN: 18 Feb 1999
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione,
 2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5S,7'aS)-
 (9CI) (CA INDEX NAME)
 OTHER NAMES:
 CN Dechloroacutamine
 FS STEREOSEARCH
 MF C19 H25 N O6
 SR CA
 LC STN Files: BIOSIS, CA, CAPLUS, TOXCENTER

Absolute stereochemistry. Rotation (-).




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

5 REFERENCES IN FILE CA (1907 TO DATE)
 5 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L8 ANSWER 15 OF 34 REGISTRY COPYRIGHT 2007 ACS on STN
 RN 121255-00-3 REGISTRY
 ED Entered STN: 23 Jun 1989
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione,
 2',3'-dihydro-3,6',7'-trimethoxy- (9CI) (CA INDEX NAME)
 OTHER CA INDEX NAMES:
 CN Acutamine, 9-dechloro-12-demethoxy-1-demethyl-11-deoxy-13-methoxy-,
 (10S,15S,16S)-
 OTHER NAMES:
 CN (+)-Limalongine
 CN Limalongine
 MF C18 H23 N O5
 SR CA
 LC STN Files: BEILSTEIN*, BIOSIS, CA, CAPLUS, NAPRALERT
 (*File contains numerically searchable property data)

Currently available stereo shown.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

1 REFERENCES IN FILE CA (1907 TO DATE)
 1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

=> file caplus
COST IN U.S. DOLLARS

FULL ESTIMATED COST

SINCE FILE	TOTAL
ENTRY	SESSION
201.35	385.83

FILE 'CAPLUS' ENTERED AT 20:11:54 ON 28 MAR 2007
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 28 Mar 2007 VOL 146 ISS 14
FILE LAST UPDATED: 27 Mar 2007 (20070327/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>

=> s 18

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom
11:Atom 12:Atom 13:CLASS 14:Atom 15:Atom 16:Atom 17:Atom

Stereo Bonds:

15-5 (Single Hash).
16-4 (Single Hash).

Stereo Chiral Centers:

4 (Parity=Odd)
5 (Parity=Odd)

Stereo RSS Sets:

Type=Relative (Default). 2 Nodes= 4 5

L10 STRUCTURE UPLOADED

=> s l10 full

REGISTRY INITIATED

Substance data SEARCH and crossover from CAS REGISTRY in progress...

Use DISPLAY HITSTR (or FHITSTR) to directly view retrieved structures.

L12 ANSWER 1 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2006:1053937 CAPLUS

DOCUMENT NUMBER: 145:460739

TITLE: Ionizing rule and characteristic spectra analysis of electrospray ionization for alkaloids in *Menispermum dauricum* DC
AUTHOR(S): Chen, Yong; Chen, Huaxia
CORPORATE SOURCE: Hubei Province Key Lab. of Bio-Technology of Traditional Chinese Medicine, Hubei University, Wuhan, 430062, Peop. Rep. China
SOURCE: Fenxi Huaxue (2006), 34(5), 675-678
CODEN: FHHHDT; ISSN: 0253-3820
PUBLISHER: Kexue Chubanshe
DOCUMENT TYPE: Journal
LANGUAGE: Chinese

AB The MS and MS2 spectra of tetrandrine and sinomenine in pos. ion detection mode were analyzed by electrospray ionization quadrupole ion trap mass spectrometry (ESI-QITMS), and their cleavage patterns were summarized. The alkaloids extracted from the medicinal materials were also analyzed


using ESI-QITMS. Tetrandrine and sinomenine were identified in the extraction by comparing the MS2 spectra of mol. ions m/z 623 and 330 with those of tetrandrine and sinomenine stds. Other known 14 ingredients were identified according to the mol. ions in MS and the characteristic product ions in MS2. Acutumine, acutumidine and acutumine, which were three kinds of new alkaloids containing chlorine found in the leaves of

Menispermum dauricum DC., were found in the extraction. The characteristic print of sixteen kinds of alkaloids (one has four kinds of isomers) in the standard medicinal materials was worked in selected ion monitor mode.

IT 17088-50-5, Acutumine 18145-26-1, Acutumidine 23512-32-5, Acutumine
RL: ANT (Analyte); ANST (Analytical study)
(anal. of alkaloids in *Menispermum dauricum* by electrospray ionization MS)

RN 17088-50-5 CAPLUS
CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5S,7'aS,9'S)- (CA INDEX NAME)

Absolute stereochemistry.



RN 18145-26-1 CAPLUS
CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-, (1R,3'aS,5S,7'aS,9'S)- (9CI) (CA INDEX NAME)

L12 ANSWER 2 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2006:1040729 CAPLUS

DOCUMENT NUMBER: 146:54913

TITLE: Aporphine alkaloids and their reversal activity of multidrug resistance (MDR) from the stems and rhizomes of *Sinomenium acutum*


AUTHOR(S): Min, Yong Deuk; Choi, Sang Un; Lee, Kang Ro
CORPORATE SOURCE: Natural Products Laboratory, College of Pharmacy, Sungkyunkwan University, Suwon, 440-746, S. Korea
SOURCE: Archives of Pharmacol Research (2006), 29(8), 627-632
CODEN: APHRDQ; ISSN: 0253-6269
PUBLISHER: Pharmaceutical Society of Korea
DOCUMENT TYPE: Journal
LANGUAGE: English

AB Chromatog. separation of the MeOH extract from the stems and rhizomes of *Sinomenium acutum* led to the isolation of nine alkaloids and a lignan. Their structures were determined to be dauriporphine (1), bianfugicine (2), dauriporphinoline (3), meniporphine (4), (-)-syringaresinol (5), N-feruloyltyramine (6), acutumine (7), dauricumine (8), sinomenine (9), and magnoflorine (10) by spectroscopic means. These compds. were examined for their P-gp mediated MDR reversal activity in human cancer cells. Compound 1 showed the most potent P-gp MDR inhibition activity with an ED50 value 0.03 μ g/mL and 0.00010 μ g/mL in the MES-SA/DX5 and HCT15 cells, resp.

IT 17088-50-5, Acutumine 345641-00-1, Dauricumine
RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(aporphine alkaloids from the stems and rhizomes of *Sinomenium acutum* and their reversal of multidrug resistance (MDR))

RN 17088-50-5 CAPLUS
CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5S,7'aS,9'S)- (CA INDEX NAME)


Absolute stereochemistry.



RN 345641-00-1 CAPLUS
CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5R,7'aS,9'S)- (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).


L12 ANSWER 1 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)
Absolute stereochemistry. Rotation (-).




RN 23512-32-5 CAPLUS

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,7'aS,9'S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L12 ANSWER 2 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)



REFERENCE COUNT: 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


DOCUMENT NUMBER: 144:103959
 TITLE: Dechlorodauricumine from cultured roots of Menispermum dauricum
 AUTHOR(S): Sugimoto, Yukihiko; Matsui, Miharui; Takikawa, Hirotsato; Sasaki, Mitsuru; Kato, Masako
 CORPORATE SOURCE: Graduate School of Science and Technology, Kobe University, Kobe, 657-8501, Japan
 SOURCE: Phytochemistry (Elsevier) (2005), 66(22), 2627-2631
 CODEN: PHYCAS; ISSN: 0031-9422
 PUBLISHER: Elsevier Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB Dechlorodauricumine, a possible organic substrate for biotransformation, was isolated from cultured roots of Menispermum dauricum, a rich source of chlorinated alkaloids. Its structure was established by spectroscopic and chemical methods.

IT 17088-50-SP, Acutumine 345641-00-1P, Dauricumine
 RL: BSU (Biological study, unclassified); NPO (Natural product occurrence); PRP (Properties); PUR (Purification or recovery); RCT (Reactant); BIOL (Biological study); OCCU (Occurrence); PREP (Preparation); RACT (Reactant or reagent)
 (alkaloid from roots of Menispermum dauricum)

RN 17088-50-5 CAPLUS
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5S,7'aS,9'S)- (CA INDEX NAME)

Absolute stereochemistry.




RN 345641-00-1 CAPLUS
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5R,7'aS,9'S)- (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).

IT 18145-26-1P, Acutumidine 345640-99-5P, Dauricumidine
 RL: BSU (Biological study, unclassified); NPO (Natural product occurrence); PUR (Purification or recovery); BIOL (Biological study); OCCU (Occurrence); PREP (Preparation)
 (alkaloid from roots of Menispermum dauricum)


RN 18145-26-1 CAPLUS
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-, (1R,3'aS,5S,7'aS,9'S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).




RN 345640-99-5 CAPLUS
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-, (1R,3'aS,5R,7'aS,9'S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).




REFERENCE COUNT: 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT



IT 219794-33-9P, Dechloroacutumine 873078-37-6P, Dechlorodauricumine
 RL: BSU (Biological study, unclassified); NPO (Natural product occurrence); PRP (Properties); PUR (Purification or recovery); SPN (Synthetic preparation); BIOL (Biological study); OCCU (Occurrence); PREP (Preparation)
 (alkaloid from roots of Menispermum dauricum)


RN 219794-33-9 CAPLUS
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5S,7'aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



RN 873078-37-6 CAPLUS
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1S,3'aS,5R,7'aS)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).




ACCESSION NUMBER: 2005:559577 CAPLUS
 DOCUMENT NUMBER: 143:226001
 TITLE: Morphinane Alkaloids with Cell Protective Effects from Sinomenium acutum
 AUTHOR(S): Bao, Guan-Hui; Qin, Guo-Wei; Wang, Rui; Tang, Xi-Can
 CORPORATE SOURCE: Shanghai Institute of Materia Medica, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences, Shanghai, 201203, Peop. Rep. China
 SOURCE: Journal of Natural Products (2005), 68(7), 1128-1130
 CODEN: JNPROD; ISSN: 0163-3864
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 143:226001

AB One new morphinane alkaloid, sinomenine N-oxide (1), and one new natural occurring morphinane alkaloid, N-demethylsinomenine (2), together with six known alkaloids, 7,8-didehydro-4-hydroxy-3,7-dimethoxymorphinan-6-ol (3), sinomenine (4), sinoacutine (5), N-norsinoacutine, acutumine, and acutumidine, were isolated from the stems of Sinomenium acutum. Their structures were elucidated on the basis of spectroscopic anal. and chemical methods. Compds. 2, 3, and 5 have protective effects against hydrogen peroxide-induced cell injury.

IT 17088-50-5P, Acutumine 18145-26-1P, Acutumidine
 RL: BSU (Biological study, unclassified); NPO (Natural product occurrence); PRP (Properties); PUR (Purification or recovery); BIOL (Biological study); OCCU (Occurrence); PREP (Preparation)
 (morphinane Alkaloids from Sinomenium acutum)


RN 17088-50-5 CAPLUS
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5S,7'aS,9'S)- (CA INDEX NAME)

Absolute stereochemistry.




RN 18145-26-1 CAPLUS
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-, (1R,3'aS,5S,7'aS,9'S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).




REFERENCE COUNT: 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L12 ANSWER 5 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2005:128007 CAPLUS
 DOCUMENT NUMBER: 142:355429
 TITLE: Synthesis of the Core Structure of Acutumine
 AUTHOR(S): Reeder, Matthew D.; Srikanth, G. S. C.; Jones, Spencer B.; Castle, Steven L.
 CORPORATE SOURCE: Department of Chemistry and Biochemistry, Brigham Young University, Provo, UT, 84602, USA
 SOURCE: Organic Letters (2005), 7(6), 1089-1092
 PUBLISHER: CODEN: ORLEF7; ISSN: 1523-7060
 DOCUMENT TYPE: American Chemical Society
 LANGUAGE: Journal
 OTHER SOURCE(S): English
 CASREACT 142:355429
 GI



AB The tricyclic core of the bioactive natural product acutumine (I) has been synthesized. Key steps include an oxidative phenolic coupling to form a masked o-benzoquinone, an anionic oxy-Cope rearrangement to construct an all-carbon quaternary center, and a Michael-type cyclization to form an amine-bearing quaternary carbon. The target compound exists in solution as an enol, in contrast to related compounds that are ketones. A model explaining these observations is presented.
 IT 17088-50-5P
 RL: PNU (Preparation, unclassified); PREP (Preparation) (synthesis of the core structure of acutumine)
 RN 17088-50-5 CAPLUS
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5S,7'aS,9'S)- (CA INDEX NAME)

Absolute stereochemistry.





REFERENCE COUNT: 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L12 ANSWER 6 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2004:2861 CAPLUS
 DOCUMENT NUMBER: 140:59819
 TITLE: Preparation and formulation of acutumine and acutumine compounds for the treatment of cognitive deficiency and neurodegenerative diseases
 INVENTOR(S): Qin, Guo-Wei; Tang, Xi-Can; Lestage, Pierre; Caigard, Daniel-Henri; Renard, Pierre
 PATENT ASSIGNEE(S): Shanghai Institute of Materia Medica, Peop. Rep. China; Les Laboratoires Servier
 SOURCE: PCT Int. Appl., 37 pp.
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004000815	A1	20031231	WO 2003-1B2600	20030616
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
CN 1465566	A	20040107	CN 2002-121479	20020625
CA 2491214	A1	20031231	CA 2003-2491214	20030616
AU 2003242278	A1	20040106	AU 2003-242278	20030616
BR 2003012444	A	20050510	BR 2003-12444	20030616
CN 1675183	A	20050928	CN 2003-819126	20030616
EP 1608625	A1	20051228	EP 2003-732907	20030616
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
JP 200601174	T	20060112	JP 2004-515147	20030616
NZ 537405	A	20060331	NZ 2003-537405	20030616
US 2006167076	A1	20060727	US 2004-519418	20041222
NO 2005000214	A	20050113	NO 2005-214	20050113
PRIORITY APPLN. INFO.:			CN 2002-121479	A 20020625
			WO 2003-1B2600	W 20030616

OTHER SOURCE(S): MARPAT 140:59819
 GI



AB Acutumine and compds. thereof of formula I [R1, R2 = H, bond; R3 = H, alkoxy; R4 = H, OH, alkoxy, alkylcarbonyloxy, arylcarbonyloxy; R5 = H, halo; R6 = H, alkyl, alkylcarbonyl, aryl; R7, R10 = alkoxy; R8R9 = bond; R8R12 = sulfide bridge; R9R10 = oxo; R13 = H, Cl; R11 = OH, alkoxy, oxo, oxime, O-alkyl oxime; R12 = H; with proviso] are prepared. The compds. can be used for the treatment of cognitive deficiencies associated with cerebral ageing and with neurodegenerative diseases. Thus, II is prepared from acutumidine, formaldehyde and propanoic anhydride. II was shown to counteract scopolamine-induced memory impairments in the Morris water maze test in mice, indicating anti-amnesic properties.

IT 23512-32-5P


RL: PAC (Pharmacological activity); PUR (Purification or recovery); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(isolation of acutumine compds. for the treatment of cognitive deficiency and neurodegenerative diseases)

RN 23512-32-5 CAPLUS

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,7'aS,9'S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 637770-93-5P 637770-94-6P 637770-95-7P
637770-96-8P 637770-97-9P 637770-98-0P
637770-99-1P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of acutumine compds. for the treatment of cognitive deficiency and neurodegenerative diseases)

RN 637770-93-5 CAPLUS

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-4,6',7'-trimethoxy-1'-methyl-5-(1-oxopropoxy)-, (1R,3'aS,5S,7'aS,9'S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.


L12 ANSWER 6 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)

RN 637770-96-8 CAPLUS

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-1'-ethyl-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-, 5'-oxime, (1R,3'aS,5S,7'aS,9'S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.


Double bond geometry unknown.



RN 637770-97-9 CAPLUS

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-4,5,6',7'-tetramethoxy-1'-methyl-, (1R,3'aS,5S,7'aS,9'S)-(9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 637770-98-0 CAPLUS

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,4',5'-trione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5S,7'aS,9'S)-(9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 637770-94-6 CAPLUS

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-1'-ethyl-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-, (1R,3'aS,5S,7'aS,9'S)-(9CI) (CA INDEX NAME)


Absolute stereochemistry.




RN 637770-95-7 CAPLUS

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-4,6',7'-trimethoxy-5-(1-oxopropoxy)-1'-(1-oxopropyl)-, (1R,3'aS,5S,7'aS,9'S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.




L12 ANSWER 6 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)



RN 637770-99-1 CAPLUS

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-6',7'-dimethoxy-1'-methyl-, (1R,3'aS,5R,7'aS,9'S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 18145-26-1, Acutumidine


RL: RCT (Reactant); RACT (Reactant or reagent)

(preparation of acutumine compds. for the treatment of cognitive deficiency and neurodegenerative diseases)

RN 18145-26-1 CAPLUS

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-, (1R,3'aS,5S,7'aS,9'S)-(9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



L12 ANSWER 6 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)

IT 17088-50-5P, Acutumine
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
(Reactant or reagent)
(preparation of acutumine compds. for the treatment of cognitive
deficiency
and neurodegenerative diseases)
RN 17088-50-5 CAPLUS
CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione,
9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-,
(1R,3'aS,5S,7'aS,9'S)- (CA INDEX NAME)



Absolute stereochemistry.



REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L12 ANSWER 7 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2002:759661 CAPLUS
DOCUMENT NUMBER: 138:103611
TITLE: Alkaloids from *Menispermum dauricum*
AUTHOR(S): Yu, Bing-Wu; Chen, Jian-Yong; Wang, Yan-Ping; Cheng,
Kin-Fai; Li, Xiao-Yu; Qin, Guo-Wei
CORPORATE SOURCE: Shanghai Institutes for Biological Sciences, Shanghai
Institute of Materia Medica, Chinese Academy of
Sciences, Shanghai, 200031, Peop. Rep. China
SOURCE: Phytochemistry (2002), 61(4), 439-442
CODEN: PHYTCA; ISSN: 0031-9422
PUBLISHER: Elsevier Science Ltd.
DOCUMENT TYPE: Journal
LANGUAGE: English
GI




AB The alkaloids, dechloroacutumidine (I) and 1-epidechloroacutumine (II),
together with three known alkaloids, acutumidine, acutumine, and
dechloroacutumine, were isolated from the rhizomes of *Menispermum dauricum*
and their structures established by spectral and chemical methods. The
cytotoxicity of each compound against the growth of human cell lines was
studied, and acutumine selectively inhibited T-cell growth.

IT 17088-50-5, Acutumine 18145-26-1, Acutumidine
219794-33-9, Dechloroacutumine
RL: BSU (Biological study, unclassified); BIOL (Biological study)
(alkaloids from *Menispermum dauricum*)

RN 17088-50-5 CAPLUS
CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione,
9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-,
(1R,3'aS,5S,7'aS,9'S)- (CA INDEX NAME)


Absolute stereochemistry.



L12 ANSWER 7 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)


RN 18145-26-1 CAPLUS
CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione,
2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5S,7'aS)-
(9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



RN 219794-33-9 CAPLUS
CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione,
2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5S,7'aS)-
(9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



IT 488736-03-4P 488736-04-5P
RL: NPO (Natural product occurrence); PRP (Properties); PUR (Purification
or recovery); BIOL (Biological study); OCCU (Occurrence); PREP
(Preparation)
(alkaloids from *Menispermum dauricum*)
RN 488736-03-4 CAPLUS
CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione,
2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-, (1S,3'aS,5S,7'aS)- (9CI) (CA
INDEX NAME)


Absolute stereochemistry. Rotation (-).

L12 ANSWER 7 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)



RN 488736-04-5 CAPLUS
CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione,
2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1S,3'aS,5R,7'aS)-
(9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L12 ANSWER 8 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2002:393345 CAPLUS
 DOCUMENT NUMBER: 138:112112
 TITLE: Nitrotyrasacutuminine from *Menispermum dauricum*
 AUTHOR(S): Yu, Bing-Wu; Chen, Jian-Yong; Zhou, Tian-Xi; Cheng, Kin-Fai; Qin, Guo-Wai
 CORPORATE SOURCE: Shanghai Institute of Materia Medica, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences, Shanghai, 200031, Peop. Rep. China
 SOURCE: Natural Product Letters (2002), 16(3), 155-159
 CODEN: NPLEEF; ISSN: 1057-5634
 PUBLISHER: Taylor & Francis Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB Nitrotyrasacutuminine, an unusual nitrated morphine-type alkaloid was isolated from the roots of *Menispermum dauricum*. Its structure was determined by various 2D spectra and chemical methods.
 IT 486429-90-7F, Nitrotyrasacutuminine
 RL: NPO (Natural product occurrence); PUR (Purification or recovery); THU (Therapeutic use); BIOL (Biological study); OCCU (Occurrence); PREP (Preparation); USES (Uses)
 (nitrotyrasacutuminine from *Menispermum dauricum*)
 RN 486429-90-7 CAPLUS
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-1-[3-(4-hydroxyphenyl)-2-nitropropyl]-4,6',7'-trimethoxy-, (1S,3'aS,5R,7'aS,9'S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).
 Currently available stereo shown.



REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L12 ANSWER 9 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2001:268660 CAPLUS
 DOCUMENT NUMBER: 135:58490
 TITLE: Chlorinated alkaloids in *Menispermum dauricum* DC. root culture
 AUTHOR(S): Sugimoto, Yukihiko; Babiker, Hind A. A.; Saisho, Tomoki; Furumoto, Toshio; Inanaga, Shinobu; Kato, Masako
 CORPORATE SOURCE: Arid Land Research Center, Tottori University, Tottori, 680-0001, Japan
 SOURCE: Journal of Organic Chemistry (2001), 66(10), 3299-3302
 CODEN: JOCEAH; ISSN: 0022-3263
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI



AB Feeding expts. using ^{36}Cl showed that *Menispermum dauricum* root culture produces four alkaloids containing chlorine. They included the novel alkaloids dauricumine (I) and dauricumidine (II) as well as the known alkaloids acutumine and acutumidine. The structures of novel alkaloids were established by spectroscopic, crystallog., and chemical methods. These four alkaloids were labeled with ^{36}Cl , isolated, and fed independently to root cultures. Mutual conversion between acutumine and acutumidine, and between dauricumine and dauricumidine by N-methylation and N-demethylation, was demonstrated. Moreover, dauricumine was converted to acutumine and acutumidine. Epimerization of acutumidine to dauricumidine or vice versa was not observed. These results suggest that dauricumine is the first chlorinated alkaloid formed in cultured *M. dauricum* roots. Skewed distribution of radioactivity derived from labeled dauricumine is proof that epimerization at C-1 proceeds at a lower rate than N-demethylation.
 IT 17088-50-5 18145-26-1
 RL: BOC (Biological occurrence); BPR (Biological process); BSU (Biological study, unclassified); MFM (Metabolic formation); BIOL (Biological study); FORM (Formation, nonpreparative); OCCU (Occurrence); PROC (Process)
 (chlorinated alkaloids in *Menispermum dauricum* root culture and study in their formation)
 RN 17088-50-5 CAPLUS
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5S,7'aS,9'S)- (CA INDEX NAME)


Absolute stereochemistry.

L12 ANSWER 9 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)




RN 18145-26-1 CAPLUS
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-, (1R,3'aS,5S,7'aS,9'S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



IT 345640-99-5P 345641-00-1P
 RL: BOC (Biological occurrence); BPR (Biological process); BSU (Biological study, unclassified); MFM (Metabolic formation); PRP (Properties); PUR (Purification or recovery); BIOL (Biological study); FORM (Formation, nonpreparative); OCCU (Occurrence); PREP (Preparation); PROC (Process)
 (chlorinated alkaloids in *Menispermum dauricum* root culture and study in their formation)
 RN 345640-99-5 CAPLUS
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-, (1R,3'aS,5R,7'aS,9'S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (+).



L12 ANSWER 9 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)

RN 345641-00-1 CAPLUS
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5R,7'aS,9'S)- (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



REFERENCE COUNT: 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L12 ANSWER 10 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1999:243783 CAPLUS

DOCUMENT NUMBER: 131:29875

TITLE: Biosynthetic relationship between acutumine and dechloroacutumine in *Menispermum dauricum* root cultures

AUTHOR(S): Babiker, Hind A. A.; Sugimoto, Yukihiro; Saisho, Tomoki; Inanaga, Shinobu; Hashimoto, Masayuki; Isogai, Akira

CORPORATE SOURCE: Arid Land Research Center, Tottori University, Tottori, 680-0001, Japan

SOURCE: *Biochemistry, Biotechnology, and Biochemistry* (1999), 63(3), 515-518

CODEN: BBBIEJ; ISSN: 0916-8451

PUBLISHER: Japan Society for Biochemistry, Biotechnology, and Agrochemistry

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The biosynthetic relationship between acutumine (I) and dechloroacutumine (II) was studied by using ¹³C-labeled tyrosine and ³H-labeled 2 as tracers. ¹³C-NMR spectra of ¹³C-labeled I and II showed that the alkaloids, each composed of two mols. of tyrosine, are derived from the same biosynthetic pathway. Feeding *Menispermum dauricum* (*Menispermaceae*) roots, cultured in a chloride-enriched medium, with ³H-labeled II demonstrated that I is the only alkaloid metabolite of II. Conversion (5%) of the exogenously applied II, taken up by the roots, into I showed that II is the precursor of I. Incomplete conversion of II into I suggests accumulation of the exogenously applied II in cell organelles and/or compartmentation of the enzymes involved in the biosynthesis of I.

IT 219794-33-9, Dechloroacutumine


RL: BPR (Biological process); BSU (Biological study, unclassified); MFM (Metabolic formation); BIOL (Biological study); FORM (Formation, nonpreparative); PROC (Process)

(Biosynthetic relationship between acutumine and dechloroacutumine in *Menispermum dauricum* root cultures)

RN 219794-33-9 CAPLUS

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole-2,5'-(4'H)-dione, 2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5S,7'aS)-(9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



IT 17088-50-5, Acutumine

RL: BSU (Biological study, unclassified); MFM (Metabolic formation); BIOL (Biological study); FORM (Formation, nonpreparative)

(Biosynthetic relationship between acutumine and dechloroacutumine in *Menispermum dauricum* root cultures)

RN 17088-50-5 CAPLUS

L12 ANSWER 11 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1999:227389 CAPLUS

DOCUMENT NUMBER: 131:4278

TITLE: Effects of chloride ion on acutumine and dechloroacutumine production by *Menispermum dauricum* root culture

AUTHOR(S): Babiker, H. A. A.; Sugimoto, Y.; Saisho, T.; Inanaga, S.

CORPORATE SOURCE: Arid Land Research Center, Tottori University, Tottori, 680-0001, Japan

SOURCE: *Phytochemistry* (1999), 50(5), 775-779

CODEN: PHYTCA; ISSN: 0031-9422

PUBLISHER: Elsevier Science Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The effects of chloride ion on the production of acutumine and dechloroacutumine, by *Menispermum dauricum* root culture, were studied. The chloride ion content in the medium plays a key role in the production of both alkaloids. A low chloride medium promoted production of dechloroacutumine and suppressed that of acutumine. Production of the two alkaloids during the 60 day culture period was closely associated with root biomass. Both alkaloids accumulated in the roots and a relatively small proportion was exuded into the medium. The intact plant produced very low amounts of both alkaloids. On the average, cultured roots contained 22- and 75-fold more acutumine and dechloroacutumine, resp., than intact plants.

IT 17088-50-5P, Acutumine 219794-33-9P, Dechloroacutumine


RL: BMF (Bioindustrial manufacture); BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PREP (Preparation); PROC (Process)

(Effects of chloride ion on acutumine and dechloroacutumine production by *Menispermum dauricum* root culture)

RN 17088-50-5 CAPLUS

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5S,7'aS,9'S)-(CA INDEX NAME)

Absolute stereochemistry.



RN 219794-33-9 CAPLUS


CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole-2,5'-(4'H)-dione, 2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5S,7'aS)-(9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).

L12 ANSWER 10 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5S,7'aS,9'S)-(CA INDEX NAME)


Absolute stereochemistry.



REFERENCE COUNT: 9

THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L12 ANSWER 11 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)



REFERENCE COUNT: 8


THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L12 ANSWER 12 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 1998:785030 CAPLUS
 DOCUMENT NUMBER: 130:122231
 TITLE: Dechloroacutumine from cultured roots of *Menispermum dauricum*
 AUTHOR(S): Sugimoto, Yukihiro; Inanaga, Shinobu; Kato, Masako; Shimizu, Toshiyuki; Hakoshima, Toshio; Isogai, Akira
 CORPORATE SOURCE: Arid Land Research Center, Tottori University, Tottori, 680-0001, Japan
 SOURCE: Phytochemistry (1998), 49(5), 1293-1297
 CODEN: PHYTCA; ISSN: 0031-9422
 PUBLISHER: Elsevier Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI



AB A novel alkaloid, dechloroacutumine (I), was isolated from *Menispermum dauricum* roots, a rich source of the chlorine-containing alkaloid acutumine, cultured in chlorine-deficient medium. Its structure was elucidated by spectral and crystallog. anal.
 IT 219794-33-9P, Dechloroacutumine
 RL: BOC (Biological occurrence); BSU (Biological study, unclassified); PRP (Properties); PUR (Purification or recovery); BIOL (Biological study); OCCU (Occurrence); PREP (Preparation)
 (dechloroacutumine from cultured roots of *Menispermum dauricum*)
 RN 219794-33-9 CAPLUS
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5S,7'aS)-(9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



REFERENCE COUNT: 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS


L12 ANSWER 13 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 1997:279419 CAPLUS
 DOCUMENT NUMBER: 127:31581
 TITLE: Effect of P-450 inhibitors on benzyloisoquinoline alkaloid biosynthesis in cultured roots of *Stephania cepharantha* and *Menispermum dauricum*
 AUTHOR(S): Sugimoto, Yukihiro; Uchida, Shinji; Inanaga, Shinobu; Isogai, Akira
 CORPORATE SOURCE: Arid Land Research Center, Tottori University, Tottori, 680, Japan
 SOURCE: Journal of Plant Physiology (1997), 150(4), 376-380
 CODEN: JPPHEY; ISSN: 0176-1617
 PUBLISHER: Fischer
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB The effect of cytochrome P 450 inhibitors was studied on biosynthesis of benzyloisoquinoline alkaloids, in cultured roots of *S. cepharantha* and *M. dauricum*. In *S. cepharantha* only 2 alkaloids, arnomoline and berbamine, were produced. Most inhibitors reduced root growth and alkaloid biosynthesis. Arnomoline and berbamine contents were pos. correlated with root growth. In *M. dauricum* ancyomidol and metyrapone promoted root growth, ketoconazole was inhibitory, while other inhibitors had inconsistent effects. Production of the alkaloids dauricine and acutumine

was curtailed by all inhibitors. Alkaloid contents were not related to root growth. None of the inhibitors induced accumulation of the immediate monomeric precursors of bis-benzyloisoquinoline. Ketoconazole-treated *M. dauricum* roots accumulated tyramine, an early precursor of benzyloisoquinoline, and 2 unidentified Tyr-derived alkaloids with mol. masses of 353 and 426.

IT 17088-50-5P, Acutumine
 RL: BOC (Biological occurrence); BPN (Biosynthetic preparation); BSU (Biological study, unclassified); BIOL (Biological study); OCCU (Occurrence); PREP (Preparation)
 (450 inhibitors effect on benzyloisoquinoline alkaloid biosynthesis in cultured roots of *Stephania cepharantha* and *Menispermum dauricum*)
 RN 17088-50-5 CAPLUS
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5S,7'aS,9'S)-(CA INDEX NAME)

Absolute stereochemistry.




L12 ANSWER 12 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L12 ANSWER 14 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 1996:198363 CAPLUS
 DOCUMENT NUMBER: 124:255828
 TITLE: Early steps of dauricine biosynthesis in cultured roots of *Menispermum dauricum*
 AUTHOR(S): Sugimoto, Yukihiro; Uchida, Shinji; Inanaga, Shinobu; Kimura, Yasuo; Hashimoto, Masayuki; Isogai, Akira
 CORPORATE SOURCE: Arid Land Research Center, Tottori University, Tottori, 680, Japan
 SOURCE: Bioscience, Biotechnology, and Biochemistry (1996), 60(3), 503-5
 CODEN: BBIEJ; ISSN: 0916-8451
 PUBLISHER: Japan Society for Bioscience, Biotechnology, and Agrochemistry
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB Cultured roots of *M. dauricum*, were fed with L-[U-14C]tyrosine, L-[3-13C]tyrosine, and [2-13C]tyramine independently, and the incorporation of possible early precursors into dauricine (I) was studied. I was composed of four mols. of tyrosine, and tyramine was specifically incorporated into the isoquinoline portions of I. Acutumine, into which 14C-labeled tyrosine was also incorporated, was identified as one of the main constituents in the alkaloid fraction from the roots.

IT 17088-50-5, Acutumine
 RL: BSU (Biological study, unclassified); MFM (Metabolic formation); BIOL (Biological study); FORM (Formation, nonpreparative)
 (formation in cultured roots of *Menispermum dauricum*)
 RN 17088-50-5 CAPLUS
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5S,7'aS,9'S)-(CA INDEX NAME)

Absolute stereochemistry.



L12 ANSWER 15 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1984:547845 CAPLUS
DOCUMENT NUMBER: 101:147845
TITLE: Isolation of (-)-stepholidine, an alkaloid of antiserotonergic-like activity from *Sinomenium acutum* Ichikawa, Kazuo; Kinoshita, Takeshi; Itai, Akiko; Itaka, Yoichi; Sankawa, Ushio
AUTHOR(S):
CORPORATE SOURCE: Fac. Pharm. Sci., Univ. Tokyo, Tokyo, 113, Japan
SOURCE: Heterocycles (1984), 22(9), 2071-7
CODEN: HETCYAM; ISSN: 0385-5414
DOCUMENT TYPE: Journal
LANGUAGE: English


AB A tetrahydropyridoberberine alkaloid, (-)-stepholidine, was isolated as an active principle showing antiserotonergic-like activity from *S. acutum* (Menispermaceae) which has been used as an oriental medicinal drug (Japanese name, Bohi; Chinese name, Fang-Ji) in Japan. An aporphine type alkaloid, liriodenine, was isolated first time from this plant along with known alkaloids hitherto obtained from this plant.

IT 17088-50-5
RL: BOC (Biological occurrence); BSU (Biological study, unclassified); BIOL (Biological study); OCCU (Occurrence) (of *Sinomenium acutum*)

RN 17088-50-5 CAPLUS

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'as,5S,7'as,9'S)- (CA INDEX NAME)

Absolute stereochemistry.



L12 ANSWER 16 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1978:500745 CAPLUS
DOCUMENT NUMBER: 89:100745
TITLE: Search for inhibitors of microorganisms among the alkaloids
AUTHOR(S): Vichkanova, S. A.; Adgina, V. V.; Izosimova, S. B.; Shipulina, L. D.; Lyutikova, L. I.
CORPORATE SOURCE: Vses. Nauchno-Issled. Inst. Lek. Rast., Bittsa, USSR
SOURCE: Khimiko-Farmatsevticheskii Zhurnal (1978), 12(2), 101-7
CODEN: KHFZAN; ISSN: 0023-1134
DOCUMENT TYPE: Journal
LANGUAGE: Russian

AB The inhibitory activity of >30 alkaloids against microorganisms and viruses was determined. Marked antimicrobial activity was observed with compds.

such as nuphleine-HCl [25249-43-8], lutenurine [63937-19-9], sanguinarine sulfate [22331-93-7], and chelerythrine [34316-15-9] whose active concns. against *Staphylococcus aureus* were 0.24-7.8 µg/mL. Most of the 30 alkaloids and alkaloid preps. tested against viruses were active, with chelidonium sulfate [66723-59-9] and O-acetylchelidonium-HCl [66723-60-2] showing particularly marked activity.


IT 66723-62-4

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study) (virucidal activity of)

RN 66723-62-4 CAPLUS

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-, hydrochloride, [3'as-[3'aa,7'aa,9'R*,10'S*(R*)]]- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



● HCl

L12 ANSWER 17 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1971:548475 CAPLUS
DOCUMENT NUMBER: 75:148475
TITLE: Alkaloids from *Menispermum canadense*
AUTHOR(S): Boskotch, Raymond W.; Knapp, Joseph E.
CORPORATE SOURCE: Coll. Pharm., Ohio State Univ., Columbus, OH, USA
SOURCE: Lloydia (1971), 34(3), 292-300
CODEN: LLOYA2; ISSN: 0024-5461
DOCUMENT TYPE: Journal
LANGUAGE: English

GI For diagram(s), see printed CA Issue.


AB Acutumine was isolated from the aerial parts of *M. canadense*, while the rhizomes yielded acutumidine, dauricine, daurinolone, N'-demethylauricine (I), magnoflorine, N-methylindocarpine methiodide (II), and dehydrocheilanthiofoline (III). The structure of the new alkaloid I was elucidated from chemical and spectral data. II and III were previously synthesized from lindcarpine and cheilanthiofoline, resp., and the isolation was the 1st report of their presence in nature.

IT 17088-50-5 18145-26-1
RL: BOC (Biological occurrence); BSU (Biological study, unclassified); BIOL (Biological study); OCCU (Occurrence) (of *Menispermum canadense*)

RN 17088-50-5 CAPLUS

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'as,5S,7'as,9'S)- (CA INDEX NAME)


Absolute stereochemistry.



RN 18145-26-1 CAPLUS

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-, (1R,3'as,5S,7'as,9'S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



L12 ANSWER 17 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)

L12 ANSWER 18 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 1971:449368 CAPLUS
DOCUMENT NUMBER: 75:49368

TITLE: Alkaloids of menispermaceae plants. CCLIX.
Alkaloids of Menispermum dauricum. Structures of
acutumine and acutumidine, chlorine-containing
alkaloids with a novel skeleton
Tomita, Masao; Okamoto, Yasuko; Kikuchi, Tohru; Osaki,
Kenji; Nishikawa, Masao; Kamiya, Kazuhide; Sasaki,
Yoshio; Matoba, Katsuhide; Goto, Kakuji
Kyoto Coll. Pharm., Kyoto, Japan
Chemical & Pharmaceutical Bulletin (1971), 19(4),
770-91
CODEN: CPBTAL; ISSN: 0009-2363

DOCUMENT TYPE:
LANGUAGE: English


GI For diagram(s), see printed CA Issue.
AB Structures of acutumine (I) and acutumidine (II), isolated from M.
dauricum DC. and Sinomenium acutum Rehd. et Wils. (Menispermaceae), were
investigated. On the basis of degradative and spectroscopic evidence,
their structures including absolute stereochemistry were assigned to the
structures I and II which agreed with the x-ray anal. These alkaloids
represent a new class of alkaloids with a novel skeleton and also provide
examples of Cl-containing alkaloids.

IT 33110-49-5P 33381-24-7P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of)

RN 33110-49-5 CAPLUS

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole-2,5'-(4'H)-dione,
9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-,
hydrobromide, [3'aS-[3'as,7'as,9'R*,10'S*(R*)]]- (9CI) (CA
INDEX NAME)

Absolute stereochemistry.



• HBr


RN 33381-24-7 CAPLUS

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole-2,5'-(4'H)-dione,
9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-,
monooxime, [3'aS-[3'as,7'as,9'R*,10'S*(R*)]]- (9CI) (CA INDEX
NAME)

CH 1


CRN 17088-50-5

L12 ANSWER 18 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)



L12 ANSWER 18 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)
CHF C19 H24 Cl N O6

Absolute stereochemistry.



CH 2

CRN 7803-49-8
CHF H3 N O

H₂N-OH


IT 17088-50-5 18145-26-1

RL: PRP (Properties)
(structure and configuration of)

RN 17088-50-5 CAPLUS

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole-2,5'-(4'H)-dione,
9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-,
(1R,3'aS,5S,7'as,9'S)- (CA INDEX NAME)

Absolute stereochemistry.



RN 18145-26-1 CAPLUS

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole-2,5'-(4'H)-dione,
9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-,
(1R,3'aS,5S,7'as,9'S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).

L12 ANSWER 19 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 1971:1113 CAPLUS
DOCUMENT NUMBER: 74:1113

TITLE: Alkaloids of Menispermaceae plants. CCLVIII.
Alkaloids of Menispermum dauricum. Basic components of
Siberian Menispermum dahuricum (Lunosemyannik
daurskii)

AUTHOR(S): Tomita, Masao; Okamoto, Yasuko; Nagai, Yoshiko;
Tanaka, Shigeko; Hayata, Toshie
CORPORATE SOURCE: Kyoto Coll. Pharm., Mukogawa Women's Univ., Kyoto,
Japan

SOURCE: Yakugaku Zasshi (1970), 90(9), 1182-6
CODEN: YKKZAJ; ISSN: 0031-6903

DOCUMENT TYPE:
LANGUAGE: Japanese

GI For diagram(s), see printed CA Issue.

AB From the rhizome of the Russian M. dauricum, the structurally known
stepharine, acutumidine, magnoflorine, and a new bisocclaurinetype base,
dauricinoline (I), were newly isolated besides dauricine, sinomenine,
acutumine, and menisperine already reported in literature. I occurs as a
pale yellow powder and its methylation with diazomethane gives
O-methyldauricine. The O,O-di-Et compound, obtained by ethylation with
diazomethane, undergoes fission by metallic Na in liquid NH₃ and produces
D-1-(p-ethoxybenzyl)-6-ethoxy-7-methoxy-2-methyl-1,2,3,4-
tetrahydroisoquinoline as the nonphenolic base and D-armepavine as the
phenolic base. From these results, the structure of I is as shown.


IT 18145-26-1

RL: BOC (Biological occurrence); BSU (Biological study, unclassified);
BIOL (Biological study); OCCU (Occurrence)
(of Menispermum dauricum)

RN 18145-26-1 CAPLUS


CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole-2,5'-(4'H)-dione,
9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-,
(1R,3'aS,5S,7'as,9'S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).




L12 ANSWER 20 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 1969:450281 CAPLUS
 DOCUMENT NUMBER: 71:50281
 TITLE: Acutumidine, new alkaloid from the leaves of Menispermum dauricum
 AUTHOR(S): Okamoto, Yasuko; Yuge, Etsuko; Nagai, Yoshiko; Katsuta, Rieko; Kishimoto, Atsuko; Kobayashi, Yoshiko; Kikuchi, Tohru; Tomita, Masao
 CORPORATE SOURCE: Fac. Pharm. Sci., Mukogawa Women's Univ., Nishinomiya, Japan
 SOURCE: Tetrahedron Letters (1969), (24), 1933-5
 CODEN: TETLEY; ISSN: 0040-4039
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI For diagram(s), see printed CA Issue.
 AB Isolation from the leaves of M. dauricum gave together with sinomenine, acutumine, disinomenine and stepharine, a small amount of a new crystalline alkaloid, acutumidine (I), m. 175-7.5°, $[\alpha]_D -100^\circ$ (CHCl₃). N.M.R., ir, uv, and mass suggested the structure for I.
 IT 23512-32-5
 RL: RCT (Reactant); RACT (Reactant or reagent) (new alkaloid from Menispermum dauricum, structure of)
 RN 23512-32-5 CAPLUS
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,7'aS,9'S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.




L12 ANSWER 21 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 1968:431127 CAPLUS
 DOCUMENT NUMBER: 69:31127
 TITLE: The x-ray analyses of acutumine and its acetate: a trial of a short cut in the structure elucidation
 AUTHOR(S): Nishikawa, Masao; Kamiya, Kazuhide; Tomita, Masao; Okamoto, Yasuko; Kikuchi, Tohru; Osaki, Kenji; Tomie, Yujiro; Nitta, Isamu; Goto, K.
 CORPORATE SOURCE: Res. Develop. Div., Takeda Chem. Ind., Ltd., Osaka, Japan
 SOURCE: Journal of the Chemical Society [Section] B: Physical Organic (1968), (6), 652-8
 CODEN: JCSPEC; ISSN: 0045-6470
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB The crystal and mol. structures of acutumine, a novel type of alkaloid containing Cl, and of its acetate, have been solved by x-ray anal. by using a three-dimensional Patterson superposition method and repeated application of least sqs. and three-dimensional Fourier methods. The result agreed with the chemical evidence obtained by concurrent degradative studies. The structure of acutumine is closely related to that of hasubanonine which was isolated from a species of the same plant family, but has a spirantype juncture of the five-membered rings A and B, with a Cl atom attached to the latter. The usefulness of applying the least-sqs. method at an unusually early stage for distinguishing real atoms from the spurious peaks appearing in the maps of min. functions or of Fourier synthesis was clearly demonstrated. In particular, observation of the behavior of temperature factors through several cycles of least-sqs. with fixed atomic coordinates was found to provide a rapid method.
 IT 17088-50-5
 RL: PRP (Properties) (structure of, calcul. of, short method for)
 RN 17088-50-5 CAPLUS
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5S,7'aS,9'S)- (CA INDEX NAME)

Absolute stereochemistry.



L12 ANSWER 22 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 1968:87430 CAPLUS
 DOCUMENT NUMBER: 68:87430
 TITLE: Structures of acutumine and acutumidine
 AUTHOR(S): Goto, Kakuiji; Tomita, Masao; Okamoto, Yasuko; Kikuchi, Tohru; Osaki, Kenji; Nishikawa, Masao; Kamiya, Kazuhide; Sasaki, Yoshio; Matoba, Katsuhide
 SOURCE: Proceedings of the Japan Academy (1967), 43(6), 499-504
 CODEN: PJACAW; ISSN: 0021-4280
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI For diagram(s), see printed CA Issue.
 AB The structures of acutumidine (I), m. 239-41° (decomposition), pKa 6.6 (50% EtOH), and $[\alpha]_D -212^\circ$ (pyridine), and acutumine (II), isolated from Sinomenium acutum or Menispermum dauricum, were established. The reduction of II acetate with LiAlH₄ gave a demethoxy dihydroxy ketone (III), m. 136-7°. Treating II with Zn in boiling Ac₂O gave a mixture of IV (R₁ = R₄ = Ac, R₂ = H, R₃ = MeO) (IVa) and IV (R₁ = R₄ = Ac, R₂ = Cl, R₃ = H) (IVb). IVa was hydrolyzed, giving a phenolic compound, which was methylated with CH₂N₂, giving IV (R₁ = R₂ = H, R₃ = MeO, R₄ = Me) (IVc). IVc was treated with Ac₂O in pyridine, giving IV (R₁ = Ac, R₂ = H, R₃ = MeO, R₄ = Me), and also oxidized with MnO₂, giving the ene-dione compound (V), m. 154-6°. The KMnO₄ oxidation of IVc gave a product, m. 75-7°, which was identified as 4,5,6-trimethoxy-1-indanone. Mild saponification of IVb, followed by methylation, gave IV (R₁ = R₃ = H, R₂ = Me), m. 219-21.5° (decomposition), which gave, on LiAlH₄ reduction followed by KMnO₄ oxidation, a compound m. 92-6°, which was thought to be 4,6-dimethoxy-1-indanone. The reduction of II with Zn-AcOH gave a dihydro compound, m. 168-71°.
 IT 17088-50-5 18145-26-1
 RL: PRP (Properties) (structure of)
 RN 17088-50-5 CAPLUS
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5S,7'aS,9'S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 18145-26-1 CAPLUS
 CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-, (1R,3'aS,5S,7'aS,9'S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).

L12 ANSWER 22 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)



L12 ANSWER 23 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 1967:517002 CAPLUS

DOCUMENT NUMBER: 67:117002

TITLE: Acutumine and acutumidine, chlorine-containing alkaloids with a novel skeleton. II. Chemical proof
AUTHOR(S): Tomita, Masao; Okamoto, Yasuko; Kikuchi, Tohru; Osaki, Kenji; Nishikawa, M.; Kamiya, Kazuhide; Sasaki, Yukio; Matoba, Katsuhide; Goto, Kakuji

CORPORATE SOURCE: Kyoto Univ., Kyoto, Japan
SOURCE: Tetrahedron Letters (1967), (25), 2425-30
CODEN: TELEAY; ISSN: 0040-4039

DOCUMENT TYPE: Journal
LANGUAGE: English

GI For diagram(s), see printed CA Issue.

AB cf. preceding abstract Based on previously reported exptl. evidence [CA 67: 54324k] and addnl. findings, 3 partial revised structures for acutumine are proposed (I, II, III). Acutumine (IV, R = Me, R' = H) (V) has a tetrasubstituted 6-membered α,β -unsatd. ketone system carrying one or two MeO groups at the α,β positions. Acutumidine acetate IV (R = Me, R' = Ac) (VI) over PtO₂ gave a dihydro compound. The N.M.R. spectrum of N,O-dibenzoylacutumidine showed a spectra consistent with the structure I. V has a secondary allyl alc. system. Treatment of VI with NaBH₄ gave a hydroxy compound (VII), showing a pos. Cotton effect at 304 m μ . The ir absorption at 1690 cm.⁻¹ and the uv band at 245 m μ in various acutumine derivs. were accordingly ascribed to a hindered five-membered conjugated ketone system. Reduction of VI with LiAlH₄ gave a dimethoxy-dihydroxy ketone (VIII), m. 136-7°, showing a pos. Cotton effect. Oxidation of V with MnO₂ gave a dehydro product (IX), ν 1745, 1695 cm.⁻¹, indicative of a 5-membered ene-dione system. The above evidence supported the partial structure II. The partial structure III followed mainly from N.M.R. spectral evidence. Treatment of V with Zn in boiling Ac₂O gave a mixture containing a neutral fraction consisting of 2 aromatic N-free products, mainly (X, R₁ = R₂ = Ac) (XI), which hydrolyzed gently to give a phenolic compound which methylated with CH₂N₂ to an O-methyl ether X (R₁ = Me, R₂ = H) (XII), and acetylated with Ac₂O-CSH₅N to the acetate X (R₁ = Me, R₂ = Ac) (XIII). The N.M.R. spectra of XI, XII, XIII indicated that the partial structure II proposed for V might remain unchanged in these products. MnO₂ oxidation of XII gave an ene-dione (XIV), m. 154-6°. XII oxidized with KMnO₄ gave a product, m. 75-7°, identified as 4,5,6-trimethoxy-1-indanone. The minor degradation product (KV, R₁ = R₂ = Ac) saponified and methylated gave an O-methyl ether XV (R₁ = Me, R₂ = H), m. 219.0-21.5° (decomposition), which reduced with LiAlH₄ and consequently oxidized with KMnO₄ to yield a small amount of crystalline 4,6-dimethoxy-1-indanone. The structure of V was

accordingly expanded to the partial structure (XVI) and the remaining moiety (C₃H₇N) was considered to form a -CH₂CH₂NMe- grouping based on N.M.R. study. Reduction of V with Zn-AcOH gave a dihydro compound (XVII),

m. 168-71°. Acetylation expts. suggested that XVII is most likely a carbinolamine, formed by cyclization of the normal reduction product (XVIII).

The circular dichroism curves of V and acutumidine IV (R = R₁ = H) have a neg. Cotton effect near 320 m μ , attributed to the $n \rightarrow \pi^*$ transition of a 6-membered α,β -unsatd. ketone system, comparable to the Cotton effect of hasubanonine (XIX) with the same structural feature of established configuration, thus establishing the structures of acutumine and acutumidine as indicated (XX, R = Me, and R = H), in accordance with structures previously obtained from x-ray analysis.


IT 17088-50-5 18145-26-1
RI: PRP (Properties)

L12 ANSWER 23 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)
(structure of)

RN 17088-50-5 CAPLUS

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5S,7'aS,9'S)-(CA INDEX NAME)


Absolute stereochemistry.



RN 18145-26-1 CAPLUS

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-, (1R,3'aS,5S,7'aS,9'S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



L12 ANSWER 24 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1967:517001 CAPLUS

DOCUMENT NUMBER: 67:117001

TITLE: Acutumine and acutumidine, chlorine-containing alkaloids with a novel skeleton. I. X-ray analysis of acutumine
AUTHOR(S): Tomita, Masao; Okamoto, Yasuko; Kikuchi, Tohru; Osaki, Kenji; Nishikawa, M.; Kamiya, Kazuhide; Sasaki, Yukio; Matoba, Katsuhide; Goto, Kakuji

CORPORATE SOURCE: Kyoto Univ., Kyoto, Japan
SOURCE: Tetrahedron Letters (1967), (25), 2421-4
CODEN: TELEAY; ISSN: 0040-4039

DOCUMENT TYPE: Journal
LANGUAGE: English

GI For diagram(s), see printed CA Issue.

AB Isolation from Sinocentrum acutum and Menispermum dauricum gave the minor alkaloid acutumine (I), m. 238-40° (decomposition), [α]_D²⁰ -206° (CSH₅N), and the N-nor base, acutumidine (II), N-Methylation of II, m. 239-41° (decomposition), [α]_D²⁰ -212° (CSH₅N), showing close similarity to I in spectral properties, gave I. A 3-dimensional sharpened Patterson function was calculated from 1459 independent data and from 14 tentative atomic positions; alternating applications of the least sqs. method and Fourier synthesis revealed the structure of I as shown (or its mirror image). Of the 14 atomic positions assumed at first, 4 were inadequate. Observed and calculated intensities of 29 pairs of reflections in the 1st and 2nd layers of Weissenberg photographs were compared from which the absolute configuration of the mol. was determined


IT 17088-50-5 18145-26-1
RI: PRP (Properties)

(structure of)

RN 17088-50-5 CAPLUS

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5S,7'aS,9'S)-(CA INDEX NAME)

Absolute stereochemistry.




RN 18145-26-1 CAPLUS

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-, (1R,3'aS,5S,7'aS,9'S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).

L12 ANSWER 24 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN (Continued)



L12 ANSWER 25 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 1967:505042 CAPLUS
 DOCUMENT NUMBER: 67:105042
 TITLE: A region of biosynthesis
 AUTHOR(S): Barton, Derek H. R.
 SOURCE: Chemistry in Britain (1967), 3(8), 330-7
 CODEN: CHMBAY; ISSN: 0009-3106
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB The developing interest in the biosynthesis of organic compds. is discussed, particularly in terms of reagent variation and its effect on specifically designed phenolic mols. The derivation of the structure of acutumine from phenolic coupling followed by further degradation is reviewed. Work previously described on the biosynthesis of sinomenine and Erythrina alkaloids (CA 67: 32851c) is also discussed. 38 references.


IT 17088-50-5

RL: PRP (Properties)
 (structure of, determination of)

RN 17088-50-5 CAPLUS

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5S,7'aS,9'S)- (CA INDEX NAME)

Absolute stereochemistry.



L12 ANSWER 26 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 1967:454324 CAPLUS
 DOCUMENT NUMBER: 67:54324
 TITLE: Characterization of acutumine
 AUTHOR(S): Goto, Kakuji; Tomita, Masao; Okamoto, Yasuko; Sasaki, Yoshio; Matoba, Katsuhide
 SOURCE: Proceedings of the Japan Academy (1966), 42(10), 1181-4
 CODEN: PJACAW; ISSN: 0021-4280
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB A revised mol. formula, C₁₉H₂₄N₂O₆Cl, is presented for acutumine (I), [pK_a 5.3, m. 238-40°, [α]_D²⁰ -206° (c 0.69, C₅H₅N)]. The presence of Cl in I was proved by mass spectra and combustion data; however, attempts to remove it by Ag₂O, LiAlH₄, or catalytic hydrogenolysis were unsuccessful. Acetylation with Ac₂O-C₅H₅N gave a monoacetylated acutumine (II), m. 162-4°. Oxidation of I with MnO₂ gave oily dehydroacutumine with formation of a new carbonyl group and the disappearance of the >CHOH proton group. I afforded a monooxime, m. 213°, which no longer shows ir absorption at 1670 cm⁻¹, indicative of one carbonyl group in I. Reduction of II with NaBH₄ gave three products, A, B, and C. A is shown by ir to have acetoxy and hydroxyl groups, and to lack the carbonyl group. Oxidation of A with MnO₂ gave II. Acetylation of A with Ac₂O-C₅H₅N afforded a diacetate, m. 149-52°. The products B and C show the absence of both the acetoxy and the carbonyl band at 1670 cm⁻¹ in the ir. Hydrogenation of II over PtO₂ in MeOH gave dihydroacutumine acetate, m. 230-1°. Treatment of II with LiAlH₄ resulted in demethoxylation and reduction of the carbonyl group to give a product m. 136-7°. Acetylation of this product with Ac₂O-C₅H₅N gave a diacetate, m. 138-42°. Oxidation of the reduction product with MnO₂ gave a dehydro product, m. 166-9°. Reaction of I with Pb(OAc)₄ gave, as a major product, a CHO-containing compound


IT 17088-50-5

RL: PRP (Properties)
 (properties of)

RN 17088-50-5 CAPLUS

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5S,7'aS,9'S)- (CA INDEX NAME)

Absolute stereochemistry.



L12 ANSWER 27 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 1961:100140 CAPLUS
 DOCUMENT NUMBER: 55:100140
 ORIGINAL REFERENCE NO.: 55:18893d-e
 TITLE: Alkaloids of Menispermum dahuricum
 AUTHOR(S): Il'inskaya, T. A.
 SOURCE: Trudy Vsesoyuz. Nauch.-issledovatel. Inst. Lekarstv. i Aromat. Rast. (1959), (No. 11), 51-64
 From: Referat. Zhur. Khim., Biol. Khim. 1961, Abstr. No. 65638.
 DOCUMENT TYPE: Journal
 LANGUAGE: Unavailable

AB Two alkaloids were isolated from M. dahuricum: sinomenine, C₁₉H₂₃O₄N m. 161-2° (EtOAc) and 180-1° (EtOH) [α]_D²⁰ -78.9° (EtOH); and acutumine, C₂₀H₂₇O₈N m. 240-1° (EtOH Et acetate 1:1) [α]_D²⁰ -120 (pyridine). Sinomenine was isolated from the total alkaloids by Et acetate recrystn. The root of the plant contained more alkaloids than any other part. All parts contained a maximum of alkaloids during bud formation and blooming.


IT 17088-50-5, Acutumine

(from Menispermum dauricum)

RN 17088-50-5 CAPLUS

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5S,7'aS,9'S)- (CA INDEX NAME)

Absolute stereochemistry.



L12 ANSWER 28 OF 28 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 1930:7549 CAPLUS
 DOCUMENT NUMBER: 24:7549
 ORIGINAL REFERENCE NO.: 24:8544-f
 TITLE: Sinomenine and disinomenine. IX. Acutumine and sinactine
 AUTHOR(S): Goto, Kakuji; Sudzuki, Hideo
 SOURCE: Bulletin of the Chemical Society of Japan (1929), 4, 220-4
 CODEN: BCSJA8; ISSN: 0009-2673
 DOCUMENT TYPE: Journal
 LANGUAGE: Unavailable

AB Cf. C. A. 24, 122. Acutumine and sinactine are 2 alkaloids recently isolated from the root of Sinomenium acutum Rehd et Wills. Acutumine has the mol. formula C₂₀H₂₇N₂O₈ or C₂₁H₂₇N₂O₈. Its absorption spectrum resembles that of narcine. It m. 199-200°. Its HCl salt shows [α]_D²⁰ 60.20°. The mol. contains 3OMe, 1CO, 1CO₂H, 1NMe and no phenolic OH groups. Sinactine m. 174°, its HCl salt decomp. at 272°, its Au double salt is amorphous; its Pt double salt, m. 245-7°. In CHCl₃, [α]_D²⁰ -312°. The mol. formula is C₁₉H₂₁N₂O₄, with 2OMe, 1 methylenedioxy, no NMe and no phenolic OH groups. The absorption spectrum almost coincides with that of laudanoline.

IT 17088-50-5P, Acutumine

RL: PREP (Preparation)

(preparation of)

RN 17088-50-5 CAPLUS

CN Spiro[3-cyclopentene-1,10']-[3a,7a]propano[1H]indole]-2,5'-(4'H)-dione, 9'-chloro-2',3'-dihydro-5-hydroxy-4,6',7'-trimethoxy-1'-methyl-, (1R,3'aS,5S,7'aS,9'S)- (CA INDEX NAME)

Absolute stereochemistry.

